FRUIT FLY CONTROL IN YOUR GARDEN IS A YEAR-ROUND ACTIVITY



START YOUR MONITORING PROGRAM 6-8 WEEKS BEFORE RIPENING BY INSTALLING TRAPS; CONTINUE MONITORING UNTIL 2-3 WEEKS AFTER PICKING.

FRUIT	AUG					SEP				ОСТ					NOV				DEC				JAN				FEB			MAR				APR				MAY				JUN				JUL				
Loquat						•																Т				П													Г											П
Apricot					(•			•													•				1													Г				Γ							
Valencia orange								•														•		•		1													Г											•
Peach																												•	•																					
Nectarine						•																							•																					
Plum					Τ					Γ															•			•	•	•																				
Feijoa Quince					Τ					Γ				Π				T					(•							•	•			•	•									
Apple - Pink Lady					T					Г				Γ				T								1		•	•						•	•	•		•	•	•	•		•						П
Grapefruit - Ruby	•		•					•										T								1														•		•			•	0	•	•	•	•
Mandarin - Imperial	•									Γ	Γ			Τ		T		T				T			\top	7														•	•						•		•	•
Navel - Leng																																									•	•	•		0	0	•	0		

NOTE: Ripening periods will vary according to fruit variety, location and season.

PRE-HARVEST

- Install monitoring traps & monitor weekly;
- If QFF are found in traps apply weekly bait sprays; or
- Install insect nets / paper bags (the earlier the better).

RIPENING AND HARVEST

- Monitor traps and check fruit for sting marks;
- Continue weekly bait sprays;
- Repair holes in nets and keep the net off the fruit;
- Destroy fallen and infested fruit (plastic bag / slash / deep burial);
- Pick fruit as it ripens and don't let it over-ripen on the tree;
- Replace/recharge monitoring trap every 3 months.

POST-HARVEST

- Continue weekly bait sprays for 2-3 weeks;
- Remove and repair nets and move to another tree; Strip pick all remaining fruit and destroy (plastic bag / slash / deep burial)